

This leaflet is a guide for physicians, nurses and pharmacists to the safe and efficient use of MabThera 1,400 mg solution for subcutaneous injection (referred to as MabThera SC).

The enclosed information pertains to the supply, storage, handling and administration of MabThera SC.

The guidance given is specific to the subcutaneous formulation only.

MabThera SC is indicated in adults for non-Hodgkin's lymphoma (NHL) only:

for the treatment of previously untreated patients with stage III-IV follicular lymphoma (FL) in combination with chemotherapy

as maintenance therapy for the treatment of patients with FL responding to induction therapy

for the treatment of patients with CD20-positive diffuse large B-cell NHL in combination with CHOP (cyclophosphamide, doxorubicin, vincristine and prednisolone) chemotherapy.

**The use of MabThera SC as monotherapy in patients with stage III-IV follicular lymphoma who are chemoresistant or are in their second or subsequent relapse after chemotherapy cannot be recommended as the safety of the once weekly administration has not been established.**

The recommended dosage is a fixed dose of 1,400 mg, irrespective of the patient's body surface area.

If you have any further questions, please refer to the full prescribing information or contact your local Roche representative.

#### Important reminder:

In case of any adverse events or safety queries please email us at [pakistan.drug\\_safety@roche.com](mailto:pakistan.drug_safety@roche.com) or call us at **+923018287010**

If you have a scientific query related to one of Roche's products, please contact us at [pakistan.medical\\_information@roche.com](mailto:pakistan.medical_information@roche.com)

Full prescribing information is attached  
**Roche Pakistan Limited**  
1st Floor, 37-B, Block 06, P.E.C.H.S.  
Karachi-75400 Pakistan  
Ph: 34538890-93, 34540731-34, Fax: 34549563



# MabThera® 1400 mg

## Solutions for subcutaneous injection (Rituximab)

### For NHL only

Note: MabThera 1400 mg is not recommended for use in CLL

Guide to supply, storage, handling and administration

Produced by Roche for healthcare professionals who are administering MabThera SC to their patients with non-Hodgkin's lymphoma (NHL).

Only for use in NHL



M-PK-00000306

## Supply, storage and handling of MabThera 1,400 mg solution for subcutaneous injection.

### How MabThera SC is supplied:

- Each carton contains one glass vial.
- Each vial contains 11.7 ml of sterile, non-pyrogenic and preservative-free solution (extractable volume is equivalent to one dose for administration to the patient).
- The solution is clear to opalescent, colourless to yellowish. Do not use if you notice unusual coloration or presence of visible particles.
- Composition
  - The active ingredient of MabThera SC is rituximab (1,400 mg per vial).
  - The excipients are:
    - Recombinant human hyaluronidase (rHuPH20): this is an enzyme used to increase the dispersion and absorption of co-administered drugs when administered subcutaneously. It allows the injection of larger volumes via the subcutaneous route.
    - Other excipients: L-histidine, L-histidine hydrochloride monohydrate, dihydrate, L-methionine, polysorbate 80 and water for injections.
  - The pH of the solution is between 5 and 6.

### How MabThera SC should be stored:

- Keep the vial in the outer carton to protect MabThera SC from light.
- Store MabThera SC in a refrigerator (2–8°C). DO NOT FREEZE.
- Check the expiry date on the outer carton.

### How to handle MabThera SC:

- Before handling MabThera SC, please check the packaging to ensure you have the correct formulation. This is in order to avoid any confusion with MabThera concentrate for solution for infusion, which has a different packaging colour code.

Check for the specific MabThera SC packaging characteristics:

1. Red labelling: **'For subcutaneous use only'**, and **'Solution for subcutaneous injection'**

2. Pink flip-off cap



- MabThera SC is ready to use; the whole content of the vial (1,400 mg rituximab) should be injected.
- MabThera SC does not contain any antimicrobial preservative and, as with all unpreserved sterile solutions, should be used immediately.

- No incompatibilities have been observed between MabThera SC and the following: polypropylene or polycarbonate syringe material, stainless steel transfer and injection needles, polyethylene Luer cone stoppers.

### How to administer MabThera SC: step-by-step guide\*

#### Important reminder:

- All patients must receive beforehand a full dose of Ristova by intravenous infusion, using Ristova intravenous formulation. The switch to MabThera subcutaneous formulation can only occur at the second or subsequent cycle of treatment.\*
- Premedication consisting of an anti-pyretic and an anti-histaminic, e.g. paracetamol and diphenhydramine, should always be given before each administration of Ristova/MabThera. Premedication with glucocorticoids should be considered if Ristova/MabThera is not given in combination with glucocorticoid-containing chemotherapy.
- MabThera SC should be administered in an environment where full resuscitation facilities are immediately available and under the close supervision of an experienced healthcare professional.

#### 1. Prepare the patient for injection

- The patient should be asked to lean back in a reclining chair or a bed and to make their abdominal region accessible for injection.



#### 2. Prepare the injection site

- The selected abdominal site should be thoroughly disinfected as per local practice.
- Each injection should be given at a different site and never into areas where the skin is red, bruised, tender, hard, or into areas where there are moles or scars.



#### 3. Prepare MabThera SC for injection

- The syringe should be prepared at the time of its administration.
- Ensure use of a needle suitable for subcutaneous injection.
- Attach the hypodermic injection needle to the syringe immediately prior to administration to avoid potential needle clogging.
- The whole content of the vial (1400 mg rituximab) should be injected.

#### 4. Perform the injection

- Pinch the skin of the abdomen with one hand to create a skin fold: this will facilitate the injection.
- Insert the injection needle into the skin fold with the other hand, using a sterile technique.
- Release the skin fold and apply pressure on the syringe, slowly injecting MabThera SC into the subcutaneous tissue.
- MabThera SC should be administered over approximately 5-7 minutes
  - Using the palm of the hand to depress the plunger can help to maintain a constant flow rate.
  - Ensure the full content of the syringe is injected into the subcutaneous tissue.
- After application, the injection site may be covered with a dressing, as per locally approved protocol.



#### 5. Inform the patient that they may leave

- The patient should be observed for at least 15 minutes following MabThera SC administration. A longer period of observation may be appropriate in patients with an increased risk of hypersensitivity reactions.
- If the patient is not receiving any further treatment after the MabThera SC injection, and if the patient is not presenting with any adverse reaction to the injection, the patient may leave the clinic.
- Many patients experience some side effects at or around the MabThera SC injection site. These local side effects include pain, swelling, induration, bleeding, skin redness, itching and rash.
- The patient should be instructed to contact their doctor immediately if the following symptoms occur: breathing difficulties, tongue or throat swelling, vomiting or palpitations, as they could be indicative of an allergic reaction.






## MabThera® 1400 mg Comparison Card for SC formulation

Please make sure that you select the correct MabThera® and Ristova® formulation & strength


**SUBCUTANEOUS INJECTION**  
For use in Non-Hodgkin's lymphoma ONLY†

MabThera 1,400 mg solution for subcutaneous injection  
Note: MabThera 1400 mg is not recommended for use in CLL



**INTRAVENOUS INFUSION**  
For use in all Ristova-approved indications†

Ristova 100 mg concentrate for solution for infusion  
Ristova 500 mg concentrate for solution for infusion



**Check for the specific MabThera SC packaging characteristics before use:**

1. Red labelling: 'For subcutaneous use only', and 'Solution for subcutaneous injection'
2. Pink flip-off cap

† Full prescribing information is attached

**Roche Pakistan Limited**  
1st floor, 37-B, Block 6, P.E.C.H.S.,  
Karachi 75400 Pakistan.  
Ph: 34538890-93, 34540731-34  
Fax: 34549563

In case of any adverse events or safety queries please email us at [pakistan.drug\\_safety@roche.com](mailto:pakistan.drug_safety@roche.com) or call us at **+923018287010**  
If you have a scientific query related to one of Roche's products, please contact us at [pakistan.medical\\_information@roche.com](mailto:pakistan.medical_information@roche.com)